




# **iCell<sup>®</sup> Hepatoblasts Prototype User's Guide**



## Trademarks

iCell is a registered trademark, and Cellular Dynamics and the  logo are trademarks of Cellular Dynamics International, Inc. and may not be used without the express written permission of Cellular Dynamics International, Inc. (CDI).

All other brands, product names, company names, trademarks, and service marks are the properties of their respective owners.

## Restrictions and Liabilities

This document is provided “as is.” CDI assumes no responsibility for any typographical, technical, or other inaccuracies in this document. CDI reserves the right to periodically change information that is contained in this document; however, CDI makes no commitment to provide any such changes, updates, enhancements, or other additions to this document to you in a timely manner or at all.

THE iCELL HEPATOBLASTS ARE PROVIDED “AS IS” TO YOU. TO THE FULLEST EXTENT PERMITTED BY APPLICABLE LAW, CDI MAKES NO, AND DISCLAIMS ALL, REPRESENTATIONS, WARRANTIES, CONDITIONS OR COVENANTS, EXPRESS OR IMPLIED, WITH RESPECT TO ANY PRODUCT REFERENCED HEREIN OR PERFORMANCE OF ANY SERVICES REFERENCED HEREIN, INCLUDING BUT NOT LIMITED TO, ANY EXPRESS OR IMPLIED WARRANTIES OR CONDITIONS OF FITNESS FOR A PARTICULAR PURPOSE, NON-INFRINGEMENT, MERCHANTABILITY, DURABILITY, TITLE, OR RELATED TO THE PERFORMANCE OR NON-PERFORMANCE OF ANY PRODUCT REFERENCED HEREIN OR PERFORMANCE OF ANY SERVICES REFERENCED HEREIN).

This document might contain references to third-party sources of information, hardware or software, products or services and/or third-party websites (collectively the “Third-Party Information”). CDI does not control, and is not responsible for, any Third-Party Information, including, without limitation the content, accuracy, copyright compliance, compatibility, performance, trustworthiness, legality, decency, links, or any other aspect of Third-Party Information. The inclusion of Third-Party Information in this document does not imply endorsement by CDI of the Third-Party Information or the third party in any way.

CDI does not in any way guarantee or represent that you will obtain satisfactory results from using iCell Hepatoblasts as described herein. You assume all risk in connection with your use of iCell Hepatoblasts.

## Conditions of Use

iCell Hepatoblasts are for life science research use only and subject to the use restrictions as contained in Appendix A. You are responsible for understanding and performing the protocols described within. CDI does not guarantee any results you may achieve. These protocols are provided as CDI’s recommendations based on its use and experience with iCell Hepatoblasts.

## Origin

iCell Hepatoblasts are manufactured in the United States of America.

## Copyright Notice

© 2016 Cellular Dynamics International, Inc. All rights reserved. This document may not be reproduced, distributed, modified or publicly displayed without the express written permission of Cellular Dynamics International, Inc.

## Revision History

Version 0.1: April 2016

# Table of Contents

<b>Before You Begin</b> .....	ii
<b>Chapter 1. Introduction</b> .....	1
Components Supplied by Cellular Dynamics .....	2
Required Equipment and Consumables.....	3
Technical Support and Training.....	3
Workflow Diagram .....	4
<b>Chapter 2. Handling and Storage</b> .....	5
Handling iCell Hepatoblasts.....	5
<b>Chapter 3. Preparing Cell Culture Surfaces</b> .....	6
Recommended Vessel: Using the Pre-coated Cell Culture Vessel.....	6
Alternative Vessel: Preparing the Collagen I Cell Culture Vessel .....	6
<b>Chapter 4. Preparing the Medium</b> .....	7
<b>Chapter 5. Thawing iCell Hepatoblasts</b> .....	8
<b>Chapter 6. Plating iCell Hepatoblasts</b> .....	9
<b>Chapter 7. Differentiating iCell Hepatoblasts</b> .....	10
Inducing Hepatocytes Differentiation .....	10
Inducing Cholangiocytes Differentiation .....	10
<b>Appendices</b> .....	11
Appendix A. Intellectual Property Rights, Use Restrictions, and Limited License...11	
Appendix B. Product Provided “AS IS” .....	11
Appendix C. Limited Liability.....	12


## Before You Begin

- Immediately transfer the frozen vials to liquid nitrogen storage.
- Read this entire iCell® Hepatoblasts Prototype User's Guide before handling or using iCell Hepatoblasts.
- iCell Hepatoblasts are for life science research use only. See Appendix A for more information and other restrictions.
- A Safety Data Sheet (SDS) for dimethyl sulfoxide (DMSO), in which iCell Hepatoblasts are frozen, is available online at [www.cellulardynamics.com/lit/](http://www.cellulardynamics.com/lit/) or on request from Cellular Dynamics International. Only technically qualified individuals experienced in handling DMSO and human biological materials should access, use, or handle iCell Hepatoblasts.

Notes

## Chapter 1. Introduction

iCell Hepatoblasts represent a progenitor population upstream in differentiation from the iCell Hepatocytes 2.0 and iCell Hepatocytes products. This cell type reflects the point in differentiation from induced pluripotent stem (iPS) cells past endoderm but before being a fully specified hepatocyte. Thus, the hepatoblast cells retain the potential to differentiate into both hepatocytes and cholangiocytes. This population of human hepatoblasts is derived from iPS cells using CDI's proprietary differentiation and purification protocols. These cells provide a reliable source of human hepatoblasts that are suitable for use in preclinical drug discovery, toxicity testing, and other life science research.



**Figure 1: iCell Hepatoblasts Represent a Population of Bipotential Cells**

*These images show iCell Hepatoblasts at day 0 post-plating and day 7 post-differentiation into hepatocytes or cholangiocytes. iCell Hepatoblasts can be induced to differentiate toward hepatocyte or cholangiocyte lineages, as demonstrated by flow cytometry, resulting in an increase of alpha-1 antitrypsin or cytokeratin-7 expressing cells, respectively.*

## Components Supplied by Cellular Dynamics

Notes

Item	Catalog Number
iCell Hepatoblasts Prototype <sup>1</sup>	HBC-100-020-001-PT
iCell Hepatoblasts Prototype User's Guide <sup>1</sup>	
Certificate of Testing <sup>2</sup>	
Certificate of Origin If required for shipping purposes	
1 Safety Data Sheets and User's Guide available online at <a href="http://www.cellulardynamics.com/lit/">www.cellulardynamics.com/lit/</a>	
2 Available online at <a href="http://www.cellulardynamics.com/cot/">www.cellulardynamics.com/cot/</a>	

## Required Equipment and Consumables

**Note:** As required for the intended use, see the following iCell Hepatoblasts Prototype Application Protocols for assay-specific equipment and consumables before thawing cells:

- Modeling Hepatocyte Differentiation with Flow Cytometry Analysis
- Modeling Cholangiocyte Differentiation with Flow Cytometry Analysis

These Application Protocols are available online at [www.cellulardynamics.com/lit/](http://www.cellulardynamics.com/lit/).

Item	Vendor	Catalog Number
<b>Equipment</b>		
37°C Water Bath	Multiple Vendors	
Biological Safety Cabinet with UV Lamp	Multiple Vendors	
Cell Culture Incubator	Multiple Vendors	
Hemocytometer or Automated Cell Counter <sup>1</sup>	Multiple Vendors	
Liquid Nitrogen Storage Unit	Multiple Vendors	
Pipettors	Multiple Vendors	
Tabletop Centrifuge	Multiple Vendors	
<b>Consumables</b>		
15 ml Centrifuge Tubes	Multiple Vendors	
Biocoat Collagen I Multiwell Plates <sup>2</sup>	Becton Dickinson	354408 (24-well) 354407 (96-well)
Pipettes	Multiple Vendors	
Trypan Blue	Life Technologies	15250
<b>Plating Medium Components</b>		
B27 Supplement	Life Technologies	17504
Blebbistatin	EMD Scientific	203391
	Sigma	B0560
BMP4	R&D Systems	314-BP
CHIR99021	Biovision	1677-25
	Stemgent	04-0004-10
FGF10	R&D Systems	345-FG
FGF basic (bFGF)	R&D Systems	233-FB
Gentamicin	Life Technologies	15750
HGF	R&D Systems	294-HGN

## Notes

Item	Vendor	Catalog Number
RPMI	Life Technologies	11875

- 1 Ensure the automated cell counter is appropriately calibrated before use.
- 2 CDI recommends using pre-coated cell culture vessels for culturing iCell Hepatoblasts. However, instructions for manually coating cell culture vessels are provided in Chapter 3, Preparing Cell Culture Surfaces.

---

## Technical Support and Training

CDI's Technical Support Scientists have the necessary laboratory and analytical experience to respond to your inquiries. In addition, in-lab training may be available upon request.


**Telephone** (877) 320-6688 (US toll-free) / (608) 310-5100 x5  
Monday - Friday, 8:30 am - 5:00 pm US Central Time

**Fax** (608) 310-5101

**Email** [support@cellulardynamics.com](mailto:support@cellulardynamics.com)

## Workflow Diagram

Notes





## Chapter 2. Handling and Storage

---

### Handling iCell Hepatoblasts

iCell Hepatoblasts are provided as cryopreserved single-cell suspensions in 1.5 ml cryovials. Upon receipt, directly transfer the individual vials containing iCell Hepatoblasts to the vapor phase of a liquid nitrogen storage dewar.

**Note:** Do not store iCell Hepatoblasts on dry ice or at  $-80^{\circ}\text{C}$  as this may impact their performance post-thawing.



It is critical to maintain cryopreserved iCell Hepatoblasts at a stable temperature. Minimize exposure of cryopreserved iCell Hepatoblasts to ambient temperature when transferring vials to liquid nitrogen storage.

## Chapter 3. Preparing Cell Culture Surfaces

iCell Hepatoblasts will plate and function optimally on collagen I-coated cell culture vessels. You can purchase pre-coated vessels (recommended) or coat them manually. Regardless of the coating methodology, have the plating surfaces ready before thawing iCell Hepatoblasts.

### Recommended Vessel: Using the Pre-coated Cell Culture Vessel

See Required Equipment and Consumables section in Chapter 1, Introduction, for ordering information for the recommended pre-coated cell culture vessels.

### Alternative Vessel: Preparing the Collagen I Cell Culture Vessel

1. Prepare 0.02 M acetic acid (Sigma, Cat. No. 320099) in sterile tissue culture grade distilled water.



Observe safety precautions for handling concentrated acids. Wear eye protection and impervious gloves. Use caution and always add acid slowly to water, never the reverse.

2. Dilute 3 mg/ml collagen I solution (Gibco, Cat. No. A1048301) in 0.02 M acetic acid to a final concentration of 100 µg/ml.
3. Select the cell culture vessel appropriate for your experimental use. Use the volumes specified in the table below in the following coating procedure. Scale volumes appropriately for other vessel formats.

Culture Vessel	Volume of 100 µg/ml Collagen I Solution (ml)	Volume of Water Rinse (ml)
24-well Cell Culture Plate	0.25	0.5
96-well Cell Culture Plate	0.032	0.1

**Table 1: Summary of Useful Volumes**

All volumes are *per well*.

4. Add collagen I solution to each well of the vessel(s).
5. Tap the vessel(s) to distribute liquid evenly.
6. Incubate the vessel(s) at room temperature for at least 1 hour.
7. After incubation, completely aspirate the collagen I solution from each well. Rinse each well 3 times with sterile tissue culture grade distilled water and aspirate completely.

**Note:** If necessary, air-dry the vessels coated with collagen I and store at 4°C for up to 7 days before use. Equilibrate the vessel(s) at room temperature before plating iCell Hepatoblasts.

## Chapter 4. Preparing the Medium

Prepare and store the Plating Medium for iCell Hepatoblasts as follows:

Component <sup>1, 2, 3, 4, 5</sup>	Amount (ml)	Final Concentration
RPMI	98	98%
B27 Supplement, 50X	2	1X
Blebbistatin, 25 mM	0.01	2.5 $\mu$ M
BMP4, 25 $\mu$ g/ml	0.2	50 ng/ml
CHIR99021, 10mM	0.03	3 $\mu$ M
FGF10, 100 $\mu$ g/ml	0.06	60 ng/ml
bFGF, 100 $\mu$ g/ml	0.01	10 ng/ml
Gentamicin, 50 mg/ml	0.05	25 $\mu$ g/ml
HGF, 25 $\mu$ g/ml	0.08	20 ng/ml

- 1 Prepare Plating Medium on the day of thaw to ensure optimal viability of cells upon plating.
- 2 Follow the manufacturer's instructions for storage and reconstitution of component stock solutions.
- 3 Filter the medium using a 150 ml, 0.2  $\mu$ m PES filter unit.
- 4 Prepare 10 ml aliquots of Plating Medium in 15 ml centrifuge tubes.
- 5 Store Plating Medium at 4°C for up to 1 week. Do not store at -20°C.

## Chapter 5. Thawing iCell Hepatoblasts

Maintain iCell Hepatoblasts in liquid nitrogen until immediately before thawing to ensure maximal performance of the cells. Complete the following steps of the thawing procedure in a time-efficient manner to facilitate optimal iCell Hepatoblasts viability and performance.

**Note:** Thaw no more than 3 vials of iCell Hepatoblasts at one time.

1. Equilibrate a 10 ml aliquot of Plating Medium in a 37°C water bath before thawing iCell Hepatoblasts.
2. Remove the iCell Hepatoblasts cryovial from the liquid nitrogen storage tank.  
**Note:** If necessary, place cryovials on dry ice for up to 10 minutes before thawing.
3. Immerse the cryovial in a 37°C water bath for 3 minutes (avoid submerging the cap) while gently swirling.
4. Immediately remove the cryovial from the water bath, spray with 70% ethanol, and place in a biological safety cabinet.
5. Gently transfer the iCell Hepatoblasts cryovial contents into the 15 ml centrifuge tube containing 10 ml of 37°C Plating Medium.



Avoid repeated pipetting of the thawed iCell Hepatoblasts cell suspension.

6. Rinse the empty iCell Hepatoblasts cryovial with 1 ml of 37°C Plating Medium to recover any residual cells from the vial. Transfer the 1 ml of Plating Medium rinse from the cryovial to the 15 ml centrifuge tube containing the iCell Hepatoblasts cell suspension. Invert the tube slowly to mix the cell suspension.
7. Centrifuge the cell suspension at 200 x g for 3 minutes at room temperature.



Avoid multiple centrifugations as this may impact viability and attachment of the cells.

8. Carefully aspirate the supernatant, taking care not to disturb the cell pellet.
9. Gently resuspend cell pellet in 5 ml of room temperature Plating Medium with a wide-bore pipette. Avoid vigorous shaking or vortexing of the cell suspension.



Avoid repeated pipetting of the thawed iCell Hepatoblasts cell suspension.

**Note:** Thaw no more than 3 vials of iCell Hepatoblasts at one time. Once thawed, pour the contents of the 2 or 3 vials into the same 50 ml centrifuge tube containing 20 or 30 ml of Plating Medium, respectively. Rinse each vial before inverting the final suspension of thawed cells.

## Chapter 6. Plating iCell Hepatoblasts

The following procedure describes how to plate iCell Hepatoblasts at  $3 \times 10^5$  viable cells/cm<sup>2</sup>. Scale volumes appropriately for other cell culture vessel formats.

1. Invert the thawed iCell Hepatoblasts cell suspension 2 - 3 times to ensure an even cell distribution before performing the cell count.
2. Remove a sample of cells to perform a cell count using a hemocytometer (using trypan blue exclusion to identify viable cells) or an automated cell counter.
3. Calculate the final volume of Plating Medium needed to obtain a desired cell plating density (i.e.  $1 \times 10^6$  viable cells/ml).
4. Dispense the cell suspension into the appropriate cell culture vessel(s). Use the volumes specified in the table below for the selected vessel format. For 24-well cell culture plates or larger, gently shake the plates to distribute the cells evenly. Do not shake 96-well cell culture plates.

Culture Vessel	Surface Area (cm <sup>2</sup> )	Plating Volume (ml)	Cell Number ( $3 \times 10^5$ cells/cm <sup>2</sup> )
24-well Cell Culture Plate	1.9	0.57	$5.7 \times 10^5$
96-well Cell Culture Plate	0.32	0.096	$0.96 \times 10^5$

**Table 2: Summary of Recommended Volumes and Measures**

*All volumes and measures are **per well**.*

5. Culture iCell Hepatoblasts in a cell culture incubator at 37°C, 5% CO<sub>2</sub> overnight.

## Chapter 7. Differentiating iCell Hepatoblasts

Notes

iCell Hepatoblasts represent a population of bipotential cells that can differentiate into hepatocytes or cholangiocytes.

---

### Inducing Hepatocytes Differentiation

iCell Hepatoblasts will differentiate into a highly pure hepatocyte population within 7 days when cultured in Hepatocyte Differentiation Medium for 1 day, then in Hepatocyte Maintenance Medium for 6 days. The differentiation can be determined and quantified as number of alpha-1 antitrypsin expressing cells.

For assay instructions, see the iCell Hepatoblasts Prototype Application Protocol: Modeling Hepatocyte Differentiation with Flow Cytometry Analysis available online at [www.cellulardynamics.com/lit/](http://www.cellulardynamics.com/lit/).

---

### Inducing Cholangiocytes Differentiation

iCell Hepatoblasts will differentiate into cholangiocytes within 7 days when cultured in Cholangiocyte Differentiation Medium containing activin A for 1 day, then in Cholangiocyte Maintenance Medium for 6 days. The differentiation can be determined and quantified as the number of cytokeratin-7 expressing cells.

For assay instructions, see the iCell Hepatoblasts Prototype Application Protocol: Modeling Cholangiocyte Differentiation with Flow Cytometry Analysis available online at [www.cellulardynamics.com/lit/](http://www.cellulardynamics.com/lit/).

## Appendices

---

### Appendix A. Intellectual Property Rights, Use Restrictions, and Limited License

A. **OWNERSHIP.** The Products are covered by pending patents and patents: <https://cellulardynamics.com/about-us/patents/>. Customer has a limited license to use the Products for internal research purposes for the sole benefit of the Customer, subject to the use restrictions included in subsection B of this Appendix A. Customer acknowledges and agrees that the receipt or purchase of the Products by Customer shall not be construed as a transfer of any title or the grant of any rights in or to the intellectual property embodied in the Products owned or licensed by Cellular Dynamics. In particular, no right or license to make, have made, offer to sell, or sell the Products, to modify or reproduce the Product or any part thereof, or to use the Products in combination with any other product(s), except product(s) provided or expressly licensed to Customer by Cellular Dynamics for such use, is implied or conveyed by the sale or transfer of Products to Customer.

B. **USE RESTRICTIONS.** The Products are licensed for internal research purposes only, and may not be used for any other purpose. The Products must be used in accordance with this User's Guide to which Customer, by ordering and accepting the Products, agrees. Customer shall not make, have made, offer to sell, or sell the Products. Customer shall not use the Products (or any modifications Customer makes to the Products or any cells derived, developed or expanded from the Products) in (i) the manufacture of any products, or (ii) any services for a third party. Customer may not transfer the Products (or any modifications Customer makes to the Products or any cells derived, developed or expanded from the Products) to any third party without Cellular Dynamics' prior written consent. Customer shall not reverse engineer the Products. Customer shall not use the Products, components or modifications thereof, or any cells derived, developed or expanded therefrom, in humans, in clinical trials, for diagnostic purposes involving human subjects, or for any investigational or other therapeutic use. Customer shall not use the Products directly or indirectly to derive or make any human gamete or gamete precursor cell. Customer shall use the Products in accordance with all applicable laws and regulations and any applicable institutional review board approved protocol and/or privacy office approval. Customer is not entitled to receive any data or information from Cellular Dynamics that directly identifies the donor of the biological materials from which the Products indirectly are derived or were made. Customer shall not attempt in any way to determine the identity of the donor of the biological materials from which the Products indirectly are derived or were made.

C. **DATA.** Customer agrees that if described on Customer's product quotation from Cellular Dynamics it will provide data and information as described therein to Cellular Dynamics regarding Customer's use of the Products.

---

### Appendix B. Product Provided "AS IS"

A. The Products are sold or provided "AS IS."

B. TO THE FULLEST EXTENT PERMITTED BY APPLICABLE LAW, CELLULAR DYNAMICS DISCLAIMS, ALL REPRESENTATIONS, AND WARRANTIES, EXPRESS OR IMPLIED (INCLUDING BUT NOT LIMITED TO, ANY IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE OR NON-INFRINGEMENT), AND ALL LIABILITY, WITH RESPECT TO THE PRODUCTS; AND BY ORDERING AND ACCEPTING THE PRODUCTS CUSTOMER WAIVES ALL RIGHTS AND REMEDIES OTHERWISE AVAILABLE WITH RESPECT TO THE PRODUCTS.

C. Customer will be solely responsible for (i) Customer's use of the Products for a purpose or in a manner other than that for which they were designed or that is permitted or in breach of the Use Restrictions above; (ii) Customer's failure to follow this User's Guide for the use, storage, and handling of the Products however such failure is caused; (iii) Customer's failure

to comply with any of the provisions of Appendix A above; and (iv) any abuse, other misuse or neglect of the Products by Customer or any damage or loss of the Products by events or occurrences beyond a person's (e.g., Cellular Dynamics') control including without limitation, accident, fire, vandalism and natural disasters (acts of God).

D. Customer acknowledges and agrees that Cellular Dynamics may fill Customer's order with any number of units of Products. Such units may be more units than Customer ordered. Customer will not be charged extra for any adjustments made by Cellular Dynamics. Because the number of cells in a unit may vary from lot to lot, Cellular Dynamics reserves the right to fill the order with that number of units which is sufficient to fill Customer's order.

---

## Appendix C. Limited Liability

TO THE FULLEST EXTENT PERMITTED UNDER APPLICABLE LAW, CELLULAR DYNAMICS SHALL NOT HAVE ANY LIABILITY FOR INCIDENTAL, COMPENSATORY, PUNITIVE, CONSEQUENTIAL, INDIRECT, SPECIAL OR OTHER SIMILAR DAMAGES, HOWEVER CAUSED AND REGARDLESS OF FORM OF ACTION WHETHER IN CONTRACT, TORT (INCLUDING NEGLIGENCE), STRICT PRODUCT LIABILITY OR OTHERWISE, EVEN IF CELLULAR DYNAMICS HAS BEEN ADVISED OF THE POSSIBILITY OF SUCH DAMAGES. NOTWITHSTANDING ANY OTHER TERM OR IMPLICATION TO THE CONTRARY, UNDER NO CIRCUMSTANCES SHALL CELLULAR DYNAMICS' LIABILITY TO CUSTOMER EXCEED THE AMOUNT PAID BY CUSTOMER FOR THE PRODUCTS TO CELLULAR DYNAMICS.

© 2016 Cellular Dynamics International, Inc. All rights reserved.